



## Abstract #14318

## Abstract Information

**Abstract ID:****14318****Abstract Title:****P-T-X evolution of the fluids associated with Li-Sn mineralization from Pedra Alta mine (Argemela, Portugal)****Authors (presenting author in bold):**

Alina Yakovenko<sup>1</sup>, **Dr. Alexandra Guedes, PhD**<sup>2</sup>, Fernando Noronha<sup>1</sup> and António Mateus<sup>3</sup>, (1)Earth Science Institute, Faculty of Sciences, University of Porto, (2)Universidade do Porto - Faculdade de Ciências, (3)Faculty of Sciences, University of Lisbon, Institute Dom Luiz (IDL)

**Abstract Text:**

Aiming the characterisation of mineral phases forming the mineralised Li-Sn quartz lodes at Pedra Alta (Permit N<sup>o</sup>1, Argemela) and of the fluids involved in the mineralising processes, a detailed petrography and mineralogy of the lodes were performed together with the study of fluid inclusions (FIs) in amblygonite, quartz, and cassiterite.

Pedra Alta is located in the Central Iberian Zone within the important Góis-Panasqueira-Argemela-Segura mineralization alignment. Is located 9 km west of the Fundão pluton within the metasediments of the Beiras Group. Tin mineralization occurs in subvertical quartz veins striking N80°E.

The studied samples are characterized by massive amblygonite, in well-developed crystals, associated with anhedral to subhedral quartz grains, displaying textured comb arrangements and mosaic aggregates. After the deposition of amblygonite and quartz, cassiterite occurs, in anhedral to euhedral grains generally zoned and include tiny phases of columbo-tantalite. The main sulphide is stannite and predates chalcopyrite. Native bismuth along with bismuthinite and silver-containing sulfosalts occur filling cracks in cassiterite. Their formation must have occurred after the development of Sn-bearing phases, as they also occur enclosed in stannite.

The petrographic examination revealed the presence of primary FIs in amblygonite, quartz and cassiterite. The study of these inclusions show that water is the main component of FIs trapped in different mineral phases. The CO<sub>2</sub>, CH<sub>4</sub>, N<sub>2</sub> and salt components are always minor. Nonetheless, fluids with a CO<sub>2</sub>-dominant volatile phase

were trapped in amblygonite and quartz, whereas fluids trapped in cassiterite show variations from CO<sub>2</sub>- to CH<sub>4</sub>-dominant volatile phase. Fls global homogenization temperatures range from: 280 °C to 315 °C for fluids in amblygonite; 310 °C to 360 °C for fluids in quartz; and 240 °C to 330 °C for fluids in cassiterite, suggesting trapping conditions at pressures higher than 50 MPa. The results indicate a temperature decrease from amblygonite to cassiterite crystallization, along with a change in fluid composition, conceivably due to changes in redox conditions and increase of rock/fluid interaction.

This work was supported by the project ERA-MIN/0003/2019 and is included in the activities of Research Group G3G of the Institute of Earth Sciences within the scope of projects UIDB/04683/2020 and UIDP/04683/2020.

**Final Paper##:**

6

**Keywords:**

Fluid inclusions, Microthermometry and Raman Microspectrometry

**Submitter's E-mail Address:**

up201503721@edu.fc.up.pt

**Submitter Full Name:**

Alina Yakovenko

**Preferred Presentation Style:**

Poster

**Conference Attendance:**

Remote

**Registration:**

I understand that the presenting author named in this submission must register for the conference by 31 May 2023.

**Submission Permission and Copyright:**

I grant non-exclusive copyright for my abstract and any related uploaded content to the Conference Exchange and the conference owners. I give permission for my details to be stored in the conference database. I, and the presenting author (if different than the submitter), agree to be contacted by the conference organizers about this submission. I agree to the recording and online streaming of any presentations scheduled as a result of this submission. I confirm all listed authors have agreed to these terms.